



## (Un)Consciousness and Time-Series Complexity A study with spontaneous EEG

Rathee, D., Cecotti, H., & Prasad, G. (2016). (Un)Consciousness and Time-Series Complexity A study with spontaneous EEG. In *Unknown Host Publication* None.

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
Unknown Host Publication

**Publication Status:**  
Published (in print/issue): 21/03/2016

**Document Version**  
Publisher's PDF, also known as Version of record

**General rights**  
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**  
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk).



# (Un)Consciousness and Time-Series Complexity

## A study with spontaneous EEG



Dheeraj Rathee, Hubert Cecotti, Girijesh Prasad

School of Computing and Intelligent Systems  
Ulster University, Derry BT48 7JL, Northern Ireland, UK




### 1. AIM

- To estimate the time-series complexity of spontaneous high-density EEG data during various levels of consciousness.


### 2. INTRODUCTION

- Awake or conscious human brains possess both *integration* of information (interactions among all regions) and *differentiation* of information (specific interaction within sub-regions). Emerging neural theories suggest reduction in these two phenomena as a strong biomarker for both level and content of consciousness.
- A recent study on EEG responses to transcranial magnetic stimulation by Casali et al [1] has shown high correlation between conscious level of brain and logarithmic complexity in electrophysiological signals. This theory-driven approach suggested a substantial decrease in the complexity of spatiotemporal pattern of triggered cortical activations with loss of consciousness.
- Another study by Schartner et al [2] studied the variations in the time-series complexity of spontaneous EEG and proposed an overall decrease in complexity of EEG signals in distinct brain regions and frequency bands.
- Here we investigate Lempel-Ziv complexity for three levels of consciousness (Awake, Mild Sedation and Moderate Sedation) with high-density spontaneous EEG data of seven subjects. The dataset was analysed earlier by Chennu et al [3] and is freely available online. Our objective is to find significant combinations of brain areas and frequency bands which demonstrate a static variation pattern of normalized complexity.

### 3. ALGORITHM



### 4. RESULTS



### 5. CONCLUSION

- We analysed the variations in time-series complexity over whole scalp and within four distinct regions. Normalized complexity indices has been generated for each regions for various frequency bands to estimate the effect of alteration of consciousness over complexity of EEG signals.
- Among all the frequency bands, only Delta( $\delta$ ) (0.5-4Hz) band shows significant reduction pattern from awake to moderate sedation condition while others fail to follow the pattern which clearly demonstrate that lower frequency EEG signals are more prominent for complexity estimation which contradicts findings of [2]. Moreover, the novel findings suggest that time-series complexity increases with increase in sedation in Alpha( $\alpha$ ) band(8-12Hz) and specifically in occipital lobe.
- Within Delta( $\delta$ ) band, except occipital lobe, all other areas show static pattern of complexity index variation.

### REFERENCES

- Casali A, Gosseries O, Rosanova M, et al. (2013) A Theoretically Based Index of Consciousness Independent of Sensory Processing and Behavior. Sci. Transl. Med. 5, 198ra105. doi: 10.1126/scitranslmed.3006294 PMID: 23946194
- Schartner M, Seth A, Noirhomme Q, Boly M, Bruno M-A, Laureys S, et al. (2015) Complexity of Multi-Dimensional Spontaneous EEG Decreases during Propofol Induced General Anaesthesia. PLoS ONE 10(8): e0133532. doi:10.1371/journal.pone.0133532
- Chennu S, O'Connor S, Adapa R, Menon DK, Bekinschtein TA (2016) Brain Connectivity Dissociates Responsiveness from Drug Exposure during Propofol-Induced Transitions of Consciousness. PLoS Comput Biol 12(1): e1004669. doi:10.1371/journal.pcbi.1004669

### ACKNOWLEDGMENT

D.R. is supported by Ulster University Vice Chancellor's Research scholarship (VCRS). G.P. and H.C. are supported by the Northern Ireland Functional Brain Mapping Facility project (1303/101154803), funded by InvestNI and the University of Ulster. G.P. is also supported by the UKIERI DST Thematic Partnership project "A BCI operated hand exoskeleton based neuro-rehabilitation system" (UKIERIDST-2013-14/126).

